




A search for missing baryons in the cosmic-web


Yan-Chuan Cai
University of Edinburgh

arXiv:1709.10378



Anna de Graaff, YC, Catherine Heymans, John A. Peacock

Baryon fraction from BBN



Fields et al. 2014
(arXiv:1412.1408)





Baryon fraction from the CMB




Baryon fraction at late-time Universe



1. Diffuse: $T < 10^5$ K, Lyman alpha absorber
2. Condensed: $T < 10^5$ K, stars and cool gas
3. Hot: $T > 10^7$ K, gas in clusters and groups
4. Warm-Hot: $10^5 < T < 10^7$ K, WHIM





Filament

125 Mpc/h

Void

Cluster


- Colberg et al. 2005MNRAS359.272C
- Clampitt et al. 2016MNRAS.457.2391C
- Epps & Hudson 2017MNRAS.468.2605E
- He et al. 2018NatAs.2.401H

Springel et al. (2005)

Thermal Sunyaev-Zel'dovich effect

CMB photons

scattered by free electrons




photon energy shifted

(Courtesy of Marian Douspis)



$$\frac{\Delta T}{T_{\text{CMB}}} = g(\nu) y$$

$$y = \int \frac{k_B T_e}{m_e c^2} n_e \sigma_T d\ell \propto n_e T_e$$




filaments and galaxy pairs

Planck SZ y-map (Planck 2015 results. XXII)




1 million pairs with:

- $6 < \text{transverse separation} < 15 \text{ Mpc/h}$
- $\text{LOS separation} < 5 \text{ Mpc/h}$
- $0.4 < z < 0.7$




SZ filaments between galaxies





Detected fSZ filament signal: $y \approx 0.6 \times 10^{-8}$ at the 5.1σ level

(Significance may change in a later revision)


Vertical profile



Horizontal profile



Vertical residual



Horizontal residual

Detected SZ filament signal: $y \approx 0.6 \times 10^{-8}$ at the 5.1σ level
(Significance may change in a later revision)

A Search for Warm/Hot Gas Filaments Between Pairs of SDSS Luminous Red Galaxies

Hideki Tanimura,^{1★} Gary Hinshaw,^{1,2,3} Ian G. McCarthy,⁴ Ludovic Van Waerbeke,^{1,2}
Yin-Zhe Ma,⁵ Alexander Mead,¹ Alireza Hojjati¹ and Tilman Tröster¹

¹*Department of Physics and Astronomy, University of British Columbia, Vancouver, BC V6T 1Z1, Canada*

²*Canadian Institute for Advanced Research, 180 Dundas St W, Toronto, ON M5G 1Z8, Canada*

³*Canada Research Chair in Observational Cosmology*

⁴*CAstrophysics Research Institute, Liverpool John Moores University, Liverpool, L3 5RF, United Kingdom*

⁵*Astrophysics and Cosmology Research Unit, School of Chemistry and Physics, University of KwaZulu-Natal, Durban, South Africa*

18 September 2017


ABSTRACT

We search the *Planck* data for a thermal Sunyaev-Zel'dovich (tSZ) signal due to gas filaments between pairs of Luminous Red Galaxies (LRG's) taken from the Sloan Digital Sky Survey Data Release 12 (SDSS/DR12). We identify $\sim 260,000$ LRG pairs in the DR12 catalog that lie within $6\text{--}10\ h^{-1}\text{Mpc}$ of each other in tangential direction and within $6\ h^{-1}\text{Mpc}$ in radial direction. We stack pairs by rotating and scaling the angular positions of each LRG so they lie on a common reference frame, then we subtract a circularly symmetric halo from each member of the pair to search for a residual signal between the pair members. We find a statistically significant (5.3σ) signal between LRG pairs in the stacked data with a magnitude $\Delta y = (1.31 \pm 0.25) \times 10^{-8}$. The uncertainty is estimated from two Monte Carlo null tests which also establish the reliability of our analysis. Assuming a simple, isothermal, cylindrical filament model of electron over-density with a radial density profile proportional to r_c/r (as determined from simulations), where r is the perpendicular distance from the cylinder axis and r_c is the core radius of the density profile, we constrain the product of over-density and filament temperature to be $\delta_c \times (T_e/10^7\text{ K}) \times (r_c/0.5h^{-1}\text{ Mpc}) = 2.7 \pm 0.5$. To our knowledge, this is the first detection of filamentary gas at over-densities typical of cosmological large-scale structure. We compare our result to the BAHAMAS suite of cosmological hydrodynamic simulations (McCarthy et al. 2017) and find a slightly lower, but marginally consistent Comptonization excess, $\Delta y = (0.84 \pm 0.24) \times 10^{-8}$.

What does the signal mean?

- I. Diffuse gas or lump structure
- II. CIB contamination
- III. Density-temperature degeneration

I. Diffuse gas or lumpy structure?



Ingredients for simulation

1. DM halo from N-body simulation;
2. Y-M relation
3. Number of galaxies per halo: HOD

II. CIB contamination





Figure 14. Expected contribution to the tSZ power spectrum for the true tSZ signal (black curve), for CIB leakage (red curve), and for tSZ-CIB leakage contribution (blue curve). The dotted line indicates a negative power spectrum.

**Planck 2015 results. XXIII. The thermal
Sunyaev-Zeldovich effect–cosmic infrared background correlation**


III. Breaking degeneracy with lensing



Data - Model = Residual

Estimated matter density for the filament: $5.6 \pm 1.6 \times \bar{\rho}(z)$

(Errorbars may change in a later revision)




Estimated matter density for the filament: $5.6 \pm 1.6 \times \bar{\rho}(z)$

Gas temperature: $T_e = 2.6 \pm 0.8 \times 10^6 \text{K}$

Account for 11% of baryons

Where are the baryons



Summary

- Detection of SZ and lensing signal between galaxies
- Lumpy haloes may contribute up to 20%
- CIB contribution is negligible
- Filament gas temperature is $\sim 10^6$ K
- Approximately 11% of baryons found