

# STUDYING THE WHIM THROUGH X-RAY EMISSION

UNIVERSITY  
OF MIAMI






Massimiliano Galeazzi


University of Miami

Enzo Branchini, Alexis Finoguenov, Anjali Gupta, Rich Kelley, Wenhao Liu, Ikuyuki Mitsuishi, Takaya Ohashi, Mauro Roncarelli, Kosuke Sato, Eugenio Ursino, Tomikkutty Velliyedathu AND the Athena WHIM Science Working Group.

# The WHIM



# WHIM X-RAY EMISSION




- Relatively Small Signal on top of background and foreground due to Galactic emission and unresolved point sources
- Grasp of current X-ray missions makes detection nearly impossible other than in a few specific cases
  - ⇒ Statistical approach using Angular Autocorrelation Function (AcF)
  - ⇒ Look in regions where a WHIM filament is expected (e.g., between clusters of galaxies)

# WHIM AcF WITH XMM-NEWTON

**AcF**

$$w(\theta) = \frac{\langle I(n)I(n') \rangle}{\langle I \rangle^2} - 1$$



**AcF of 6 XMM-Newton empty fields  
in the 0.4-0.6 keV range**




**The fraction of X-rays due to the  
WHIM  
in the energy range 0.4-0.6 keV  
is  $12 \pm 5\%$  of the total diffuse X-ray  
emission**

# The AcF in the CDFS with XMM-Newton

- 4 MS
- 33 pointings
- 2 different sets of roll-angle (180 deg apart)
- Chandra-detected point sources
- Diffuse sources from Alexis Finoguenov
- Diffuse emission consistent with other work



# The AcF in the CDFS with XMM-Newton



➤ Small correlation signal detected in both 0.4-0.6 keV and 0.7-0.9 keV band

➤ **We ruled out a systematic origin of the signal**


➤ **Negligible correlation at high energy**

➤ **The signal in the two soft bands is correlated → same origin**

➤  **$T \sim 9 \times 10^6$  K**



# The AcF in the CDFS with XMM-Newton



➤ Small correlation signal detected in both 0.4-0.6 keV and 0.7-0.9 keV band


➤ **We ruled out a systematic origin of the signal**

➤ **Negligible correlation at high energy**

➤ **The signal in the two soft bands is correlated → same origin**

➤  **$T \sim 9 \times 10^6$  K**

# The AcF in the CDFS with XMM-Newton



➤ Small correlation signal detected in both 0.4-0.6 keV and 0.7-0.9 keV band

➤ We ruled out a systematic origin of the signal



➤ Negligible correlation at high energy

➤ The signal in the two soft bands is correlated → same origin

➤  $T \sim 9 \times 10^6$  K



# Comparison with models




Signal in 0.4-0.6 keV band inconsistent with models

- Cosmic variance (it is known the the CDFS is devoid of significant structures)
- Problem with the models
  - Metal diffusion? (but what about UV data?)
  - Temperature distribution?
  - ???

# SEARCH FOR INTERCLUSTER FILAMENTS

## A222-A223




6.02e-08 1.80e-07 4.22e-07 9.00e-07 1.87e-06 3.78e-06 7.58e-06 1.53e-05 3.04e-05

- $z=0.21$
- identified as a possible filament through weak lensing and as an overdensity of color selected galaxies
- The clusters are separated by  $14'$  (2.8 Mpc)
- Projected distance 15 Mpc
- 144 ks pointing with XMM-Newton

- Filament outside the virial radius of the clusters
- emission measure  $(n_e \times n_p \times V) = 1.72 \times 10^{65} \text{ cm}^{-3}$
- Temperature:  $kT = 0.91 \text{ keV}$
- Metallicity of 0.2 solar


# SEARCH FOR INTERCLUSTER FILAMENTS

## Sculptor Supercluster



*K. Sato et al., 2010*


## Shapley Supercluster




*I. Mitsuishi et al., 2012*

# X-RAY EMISSION FROM INTERCLUSTER FILAMENTS



## Sculptor Supercluster



## Shapley Supercluster




# X-RAY EMISSION FROM INTERCLUSTER FILAMENTS





# SEARCH FOR INTERCLUSTER FILAMENTS

## Shapley point sources



**Excess emission due to galaxies in the filament between the two clusters!**



# What can we do next?

---

## 1) SHORT TERM:

Look at other deep fields with XMM-Newton (e.g., COSMOS – in progress)

## 2) SHORT-MEDIUM TERM:

Use other tracers to highlight the x-ray signal

- Cross-correlation with galaxy maps
- Cross-correlation with SZ

## 3) MEDIUM-LONG TERM:

High energy resolution studies with Athena



# The WHIM with Athena




Figure 1: Simulated emission and absorption line spectra captured in a single *Athena+* observation for two filaments at different redshifts. Lower panel: absorption spectrum from a sight line where two different filamentary systems are illuminated by a bright background source. Upper panel: corresponding emission from a 2'x2' region from the same filaments for 1 Ms exposure time. The high spectral resolution allows us to distinguish both components. *Athena+* will be able to study dozens of these sight lines in detail.

# The WHIM with Athena




Figure 3: Same line statistics as in Fig. 2 but referring to filaments detected in emission as a function of line surface brightness (Takei et al. 2011). We expect to detect serendipitously about 4 filaments in the field of view of X-IFU for observations longer than 100 ks. We have conservatively assumed that for about 30% of the systems detected in absorption, the associated X-ray line emission will be detected. The upper band illustrates the scatter in model predictions and refers to simultaneous detections of O VII and O VIII from all the baryons. The lower band refers to line emission from baryons in filaments only. This component increasingly dominates low SB. For observations longer than 100 ks, we expect that about 50% of the filaments detected are due to the WHIM.

# Future plans

- **Cross correlation with galaxies**
- **Multi-wavelength study (correlation with other bands)**
- **Extend XMM-Newton work to investigate cosmic variance**

**Table 1.** Summary of XMM-Newton Deep and Extended targets that will be used in the investigation.

Target	Area	Total Exposure	Avg Exposure	Sensitivity (0.5-2.0 keV)
	deg <sup>2</sup>	Ms	ks	10 <sup>-15</sup> erg s <sup>-1</sup> cm <sup>-2</sup>
XMM-CDFS <sup>1</sup>	0.25	3.45	3.45	0.66 (2-10 keV)
XMM-XLL <sup>1</sup>	50*	6.9	~10	5
XMM-Cosmos <sup>2</sup>	2	1.5	~68	1.7
XMM-SStripe 82 <sup>3,4</sup>	20	~0.8	4.5	5
Lockman Hole <sup>5,6</sup>		~0.7	~700	1.9

<sup>1</sup> Pierre et al., 2015, A&A, in press, D.O.I. 10.1051/0004-6361/201526766

<sup>2</sup> Cappelluti et al., 2009, A&A, 497, 635

<sup>3</sup> LaMassa et al., 2013, MNRAS, 436, 3581

<sup>4</sup> XMM AO13 proposal summary


<sup>5</sup> Worsley et al., 2004, MNRAS, 352, L28

<sup>6</sup> Brunner et al., 2008, A&A, 479, 283

<sup>7</sup> Ranalli et al., 2013, A&A, 555, A42

\* The XMM-XLL survey covers two fields of 25 deg<sup>2</sup> each. One field is in the northern galactic hemisphere, roughly at the anticenter position. The second field is in the southern hemisphere close to the edge of the galactic bulge.

# Future plans



Multi-wavelength coverage and sensitivity in the COSMOS field

# WHIM in emission. COSMOS


## Spatial information

Correlation with galaxies can beat down the foreground

$$\sigma_{\text{line}} = \frac{f_{\text{line}}}{\sqrt{[f_{\text{line}}] + (f_{\text{CXB}} + f_{\text{G}} + f_{\text{inst}})\Delta E}} \sqrt{\Delta\Omega t_{\text{exp}} A_{\text{eff}}},$$

20K galaxies in the ZCOSMOS survey  
(+20K with photometric redshifts) to trace  
the galaxy density field up to  $z=1$ .  
(Kovac+ 2010).


Several large scale structures already  
detected. Most notably at  $z\sim 0.4$  and  
at  $z\sim 0.7$ . For the latter, Wall-like, a dedicated  
redshift survey of galaxies galaxies has been  
done to increase sampling.



# WHIM in emission. Alternative probes

Probes different from galaxies are being used to statistically detect the WHIM signal. And could be combined with similar results obtained by Athena.

One example is the SZ-signal. An indirect, statistical WHIM detection has recently been claimed by de Graaff+ 2017 in filaments connecting galaxy pairs as a result of a stacking analysis.



A similar procedure can be repeated in the COSMOS field observed by Athena to look for a OVII/OVIII emission enhancement in between galaxy pairs. The resulting signal can be combined with the analogous SZ enhancement.

# X-RAY/SZ CORRELATION

Sunyaev-Zel'dovich effect


$$\frac{\Delta T_{CMB}}{T_{CMB}} = g(x)y$$

$$y = \int \sigma_T n_e \frac{k_B T_e}{m_e c^2} dr$$

$$g(x) = x \coth\left(\frac{x}{2}\right) - 4, \quad \text{with} \quad x = \frac{h\nu}{k_B T_{CMB}}$$

$$F(X\text{-rays}) \propto n_e^2$$
$$\Delta T_{SZ} \propto n_e^{1.2}$$

→ X-rays and SZ can probe different regions of the WHIM







# X-RAY/SZ CORRELATION

Simulated signal and background for 150 GHz ACT band  
and 100 ks XMM-Newton observation

## SZ signal alone

## SZ/X-ray cross coefficient



WHIM

BACKGROUND